

COMPLEXITY AND DYNAMICS

 **sidestone**press



This is a free offprint – as with all our publications the entire book is freely accessible on our website, and is available in print or as PDF e-book.

www.sidestone.com

COMPLEXITY AND DYNAMICS

*Settlement and landscape from the Bronze
Age to the Renaissance in the Nordic
Countries (1700 BC–AD 1600)*

MARIE **O**DEGAARD AND INGRID **Y**STGAARD (EDS)

© 2023 Individual authors

Published by Sidestone Press, Leiden
www.sidestone.com

Imprint: Sidestone Press Academics
This book has been peer-reviewed. For more information see www.sidestone.com

Lay-out & cover design: Sidestone Press
Photograph cover: Annette Græsli Øvrelid, Museum of Archaeology, University of Stavanger. Back cover: Magne Samdal, Museum of Cultural History, University of Oslo

ISBN 978-94-6427-042-6 (softcover)
ISBN 978-94-6427-043-3 (hardcover)
ISBN 978-94-6427-044-0 (PDF e-book)

Contents

Foreword	7
List of contributors	9
Settlement organization in Iron Age Scandinavia and beyond: Traditions, terminologies, regionalities and methodologies	11
Marie Ødegaard and Ingrid Ystgaard	
Early Iron Age village formation in Jutland, Denmark	29
Niels Haue	
Settlement structure and landscape use in Southwest Norway in the last millennium BC	41
Trond Meling	
Madla Sør in Rogaland, Southwest Norway – a settlement with long continuity?	53
Satu H. Lindell	
Maintaining boundaries: Early Iron Age settlement dynamics and spatial organisation at Dilling in Southeast Norway	69
Marie Ødegaard, Lars Erik Gjerpe and Linnea Syversætre Johannessen	
“The 207 BC Dust Veil Event” and the advent of iron reaping tools in Scandinavia	85
Lars Erik Gjerpe	
Uncovering population dynamics in Southeast Norway from 1300 BC to AD 800 using summed radiocarbon probability distributions	99
Kjetil Loftsgarden and Steinar Solheim	
Plant remains as sources to cultural history in Southeast Norway	113
Karoline Kjesrud, Luka Natassja Olsen, Irene Teixidor-Toneu, Jade J. Sandstedt, Anneleen Kool and Linda Christiansen	
Social dynamics at the Augland ceramic workshop: The introduction of soapstone in paste recipes from the Roman Iron Age and early Migration Period in southernmost Norway	129
Christian Løchsen Rødsrud and Per Ditlef Fredriksen	

Activities and community organization in Roman Iron Age Vik, Ørland, Central Norway	143
Ingrid Ystgaard	
Old Uppsala, Eastern Sweden: Framing an Iron Age tributary society	157
Per Frölund	
Burial – settlement relations at Forsandmoen, Southwest Norway	171
Barbro I. Dahl	
The elite settlement at Ströja, Eastern Sweden, AD 450–1000	187
Björn Hjulström and Marta Lindeberg	
Development of the medieval villages in Southern Finland	201
Tuuli Heinonen	
Anebjerg - A rural farm from the 17th century: Considerations on Renaissance building traditions in Denmark and what can be expected beyond the Middle Ages	211
Louise Søndergaard	

Plant remains as sources to cultural history in Southeast Norway

Karoline Kjesrud, Luka Natassja Olsen,
Irene Teixidor-Toneu, Jade J. Sandstedt,
Anneleen Kool and Linda Christiansen

Abstract

Plant remains are valuable sources for cultural history. Humans and animals live inextricably together with plants. This article investigates how a large dataset of botanical macrofossils can give insights into cultural history in southeastern Norway in the period 400 BC–AD 400. In this period, iron was introduced in the production of weapons and tools, which led people to change practices in their interaction with nature. Farming is dependent on a great variety of environmental resources for cultivation and the gathering of food, medicine, and fodder. By combining archaeological and botanical data from 40 localities from Viken and Innlandet counties this article investigates the macrofossil diversity within the localities, as well as the regions they belong to. All archaeobotanical finds from the localities have been systematized and analyzed quantitatively. The results reveal patterns in regional biodiversity as well as a representation of species in structures and features in archaeological sites.

Keywords: Archaeobotany, plant history, plant utilization, South-eastern Norway, regional variances

Introduction

Plants are and have always been essential for human life and activities; this is as true for cereals and timber as well as other plants used as food, medicine, or material culture. Archaeobotanical remains from archaeological excavations serve as important sources for interpreting past practices in farming, foraging, food, fodder, medicine, and rituals (Day 2013; Jacomet 2013; van der Veen 2018). Such “ecofacts” may provide a bridge to past concepts of nature and human-nature relations (Richer and Gearey 2018) because they are actual remains of, and sources to the study of, landscape and vegetation and therewith provide concrete insights about the environment that people perceived. This article explores how plant remains sampled from 40 localities in Innlandet and Viken counties in southwestern Norway can yield nuanced information on how people interacted with plants in the period 400 BC–AD 400.¹

Large-scale analyses of archaeobotanical remains may contribute to an awareness about typological patterns in the representation of macrofossils from archaeological excavations and hence be helpful for future analyses of more detailed plant utilization. Combined results from all sites are expected to display great variance when it comes to the representation of species, the actual number of macrofossils and their archaeological relation. In this article,

we investigate the following three areas: 1) if there are observable patterns in the distribution of archaeobotanical remains within site types, structures, and features from the selected area and time period, 2) the species richness within the localities, and 3) whether historical regional variance in macrofossil diversity in Innlandet and Viken may be revealed through big data analyses.

The results from these investigations serve as a foundation for discussing how archaeobotanical results may be used for exploring further cultural historical activity in the localities. Through these investigations, we also identify some ever-present challenges in the sampling strategies and pinpoint information that could provide more detailed insights into past human-plant interference.²

Background

Much of the historical interest in studying people's interference with plants centers around the domestication and cultivation of land, especially in the Iron Age, since this period is characterized by important innovations in farming and cultivation (Solberg 2003 [2000]; Myhre 2002; Welinder *et al.* 2004). The interest in studying technological inventions and abilities of crop domestication can be said to be founded on an economic view of nature – nature being a supply of resources – going back to the 17th century (Hverven 2018). People's understanding of the concept of nature has undergone drastic changes throughout history in close relation to changes in ideological and religious convictions (Taylor 2020). People relate to nature, and they do this with both conscious and subconscious intentions and motivations. In a semiotic mindset (Lotman 2008 [1996]), ecofacts and human messages of various kinds are all important jigsaw puzzle pieces in the bigger picture of past societal organization, cultural encounters, technological development, cognitive status, climatic changes, and environmental impact. For example, Norwegian toponyms preserve insights regarding how places have been used or how the surrounding landscape has been perceived.

In Norway, pollen analyses (microfossils) have been widely integrated as a source to vegetation history (Fægri 1944; Hjelle 2005; Høeg *et al.* 2019), whereas macrofossils are traditionally less used in analyses of historical perspectives on human-environmental interactions. Pollen diagrams can illustrate changes in species composition and communicate how plants and forests have formed the landscape from before the period of ice melting up to today through processes such as deforestation, ground clearing, and climate change, as well as other landscape changes caused by human activity, such as fodder harvesting and pasturing (Høeg 1996:11–14). The different plant species' pollination strategies influence the abundance of pollen in a site, which biases the understanding of the cultural implications in the landscape. Moreover, the diagrams often focus on the plants as indicators of various anthropogenic activities in

the landscape and do not necessarily consider the implicit value of the plant and therewith their cultural uses (Richer and Gearey 2017). The taxonomic resolution is typically lower for pollen identification than for macrofossils, which makes it easier to link pollen to a plant family, whereas the seeds often provide more information for determining the plant species.

Archaeobotanical sampling (macrobotanicals) has been integrated into regular excavation practice in Norway since the 1980s (Hjelle *et al.* 2017 with references). Macrobotanicals enable studies of the activity within a location since soil samples can be compared from different activity areas within a site. Suitable methods for interpreting such data have been outlined focusing specifically on cereals and grain cultivation (Grabowski 2014). The amount, quantity, and analytical treatment of excavated soil-samples vary from site to site. Possible use areas connected to corresponding plants are occasionally suggested in the archaeological reports, often with reference to more recent knowledge of plant traditions (e.g., Høeg 1996:151–153).

There are many unknown facets of people's interaction with plants in the Iron Age. Increased population growth of both people and their domestic animals in the Iron Age required more food and fodder (Myhre 2002; see also Gjerpe this volume; Loftsgarden and Solheim this volume). As in more recent times, many of the harvested plants were probably gathered in the outfields (Teixidor-Toneu *et al.* 2020) and represented a significant contribution to past farming societies (e.g., Bharucha and Pretty 2010; Turner *et al.* 2011). Since broad data sampling is usually the standard in all excavations, plant history can be extracted from these excavations, even without being the initial aim and purpose of the excavation.

Materials and methods

The dataset for this study consists of reported finds of macrofossils from excavations performed in southeastern Norway in the period 1993–2018 by the Museum of Cultural History, University of Oslo (KHM).³ The archaeological period ranges from 400 BC–AD 400 and includes macrofossils collected from 40 localities in the former counties of Østfold and Akershus (since 2020, part of the new county Viken), Hedmark and Oppland (since 2020, united in the new county Innlandet).⁴ Innlandet is represented by 14 localities with 18 farm numbers. Viken is represented by 26 localities with 30 farm numbers. Larger excavations sometimes contain more than one locality (with multiple farm numbers). These are here combined.

The macrofossil samples were analyzed by external laboratories post-excavation. The laboratories are based in Nordic countries such as Norway, Sweden, and Denmark. In the current dataset, macrofossil data is compiled from the lab reports themselves or excerpts from these reports (which are presented in archaeological reports).




Figure 1. Number of reported archaeobotanical finds per site type (a), structure (b) and feature type (c). NA means “Not Available”.

For this study, we included all macrobotanicals that we could access information on, both dated and undated (primary sources for dataset is listed in the bibliography). Microfossils (pollen) are not included. The majority of microfossils sampled by KHM today are not dated, but context material from the same feature is often dated and is often used as a period indicator for the microfossils. Charred material, dominated by cereals and charcoal, is often prioritized for analyses. Charcoal and wooden remains are particularly valuable since they provide an opportunity to put a date on the use and place in question (ongoing ^{14}C -database construction at KHM⁵). Among seeds, cereals are considered valuable since they are larger and thus provide more material for dating analyses compared to smaller wild seeds, but also because cereals indicate cultivation and domestication of land (Hjelle *et al.* 2017:303). Few uncharred seeds and plant remains are likely to be old, unless there are particular preservation conditions (waterlogging or mineralization) in the excavation area (Sture 2016 with references). Wild herbs have very few dated representatives. Consequently, the dataset consists of a majority of undated macrobotanicals, relying on the dating of context material. By presenting the data side by side, the possibilities and the challenges in the material can be addressed independent of dating.

For the investigation of observable patterns in the distribution of archaeobotanical remains (the first of the areas of investigation listed in the Introduction to this paper 1), the data is organized through a set of parameters. The two counties of Viken and Innlandet, containing 25 municipalities, were chosen because they are known to contain localities with large macrofossil quantities. A locality contains one or more site type (e.g., settlements, production sites). Within a site type there are one or more structures, such as houses, graves etc.,

whereas features in a site type are typically postholes, forges, cooking pits etc. Sometimes processed plant materials are found within features. These objects, such as wooden plugs, textile cloths etc., may be species determined, or dated. The macrobotanicals are sometimes recognized as plant parts such as seeds, bark, charcoal, nuts, or fruit which may or may not be botanically identified at a species level. The preservation status may be charred, dried, waterlogged, or mineralized – however this is not always stated in analysis reports.

The site types were classified into ten categories, structures were grouped into six categories and features were grouped into ten (fig. 1). Plant taxa appearances in specific site types, structures, and features were explored using a generalized mixed-effect regression model (family binomial) with locality (i.e. excavation site) as a random effect. This model allows analyzing non-normal data such as counts and percentages when random effects are present. Here, random effects are used to avoid pseudo-replication (i.e. repeated measures from the same location; Bolker *et al.* 2009).

For the investigation of the distribution of species richness within the localities (the second of the areas of investigation listed in the Introduction to this paper), macrobotanicals have been systematized in groups of tree, cereal and herbs, and macrofossil data was systematized in spreadsheets for the two counties (tab.1 and 2). The column *Total taxa no.* refers to the total number of species (taxa diversity) found at a given site and is represented by macrofossils such as charcoal, seeds, bark, buds, stems, nuts, cone scale, and leaves. *No. of tree taxa* represents the total number of tree taxa represented by nuts, charcoal, leaves, or bark. *No. of herb taxa* represents herbaceous plant taxa determined from seeds. Herbaceous plants do not have woody stems and many of them may be regarded

Locality	Farm nr.	Site type	Total no. of taxa	No. of tree taxa	No. of cereal taxa	No. of herb taxa	Total no. of macrofossils	Qu. of tree macrofossils (of which fruit/leaf/bark)	Qu. of cereal macrofossils	Qu. of herb macrofossils	Taxa of highest abundance in a feature/ Taxa of second highest abundance in a feature	Quantity (dated)	Structure	Feature
Leikvang	14/2	settlement	21	10	2	9	599	555**	9	35	Galium spurium Hordeum vulgare	8 (365-425 AD) 4 (35-30 AD)	longhouse longhouse	posthole posthole
Ljøstad	296/1, 2	settlement	18	7	4	7	340	298** (1)	34**	8**	Cerealia Brassica sp.	2 (45-250 AD) 1 (215-420 AD)	house area house area	posthole posthole
Vevla	26/1	settlement	13	8	1	4	242	232** (1)	5	5	Persicaria lapathifolia Cerealia	2 1 (0-85 AD)	longhouse longhouse	posthole posthole
Void	6/1	settlement	21	11	3	7	1296	1278** (59)	4	14	Chenopodium album Galium aparine	2 7	cultivation layer cultivation layer	layer 3 layer 3
Aker	7/201	settlement	50	8	3	39	4533	141 (42)	2787	1605	Stellaria media Stellaria media	197 192	house area house area	posthole posthole
Berg	138/1, 4	settlement/ cultivation traces	9	4	2	3	56	39**	5	12	Spergula arvensis Hordeum vulgare*	9 1 (250-375 AD)	house area house area	posthole posthole
Fulu store	3/8, 10	settlement	18	12	5	1	329	297** (4)	30**	2	Hordeum vulgare var. Nudum Triticum turgidum subsp.	19 (405-375 BC) 3	- -	burial / drying pit burial / drying pit
Meistrøm	18/9	settlement	10	7	3	0	44	39** (1)	5	0	Cerealia Secale cereale	3 1	- -	burial burial
Nyhuset Haukstad	94/3060	settlement	13	4	3	6	604	272	224	108	Avena sativa Stellaria media	70 39	- -	bloomery furnace bloomery furnace
Valum gård	20/1	settlement	38	4	5	29	2473	12**	889	1572	Chenopodium album Hordeum vulgare var. vulgare	568 388	longhouse house	posthole -
Trogstad	16/1, 17/1	settlement/ cultivation layer	22	15	3	4	271	249** (2)	19	3	Cerealia Hordeum vulgare	9 2 (680-890 AD)	house house	forge/smithy posthole
Børstad	4/1	cooking pit locality/ production site (iron)	11	8	2	1	222	207**	4	11**	Linum usitatissimum Avena sativa	10 (350-110 BC) 2	- -	forge forge
Hveem	72/1	settlement/ cultivation traces	15	4	2	9	108	87	9	12	Hordeum vulgare Persicaria sp.	6 (200-125 BC) 1 (15-90 AD)	- -	cultivation layer posthole
Døhlen	66/6	settlement/ cultivation traces	29	2	4	23	1272	355 (4)	420	497	Chenopodium album Cerealia	146 (180-60 BC) 192	house area house area	posthole posthole
Sum							12389	4061	4444	3865				

Table 1. Table presenting the excavated locations from Innlandet and the reported macrofossils from each locality organized in columns presenting the total number of taxa, divided in the number of tree taxa, cereal taxa and herb taxa from each site, as well as the total number of macrofossils and how they are distributed between trees, cereal and herbs.

** value is set to 1 where quantity was not given. Fragments are counted as 1 seed. *when more than one species have the same highest quantity in a locality the example is chosen at random. Quantity is given along with available dating results. If context material has been dated (material found together with macrofossil in the same feature), this is indicated in italics. Note that "seeds of *Hordeum vulgare*" in Valum gård was given for the whole structure and not per feature.




Figure 2. A dendrogram showing clusters of excavated localities based on similarities of species presence/absence.

Locality	Farm nr.	Site type	Total no. of taxa	No. of tree taxa	No. of cereal taxa	No. of herb taxa	Total no. of macrofossils	Qu. of tree macrofossils (of which fruit/leaf/bark)	Qu. of cereal macrofossils	Qu. of herb macrofossils	Taxa of highest abundance in a feature/ Taxa of second highest abundance in a feature	Quantity (dated)	Structure	Feature
Borgen	30/1	settlement	6	5	1	0	650	646	4	0	Cerealia	2 (260-450 AD)	longhouse	posthole
Brustad	19/1	settlement / cooking pits production site (iron)	16	8	3	5	375	365 (1)	4	6	Cerealia Papaver sp.* Corylus avellana	1 (950-530 AD) 1 1 (130-230 AD)	longhouse longhouse longhouse	posthole posthole posthole
Bråte	78/11	settlement	26	10	4	12	853	816	9	28	Cerealia Persicaria lapathifolia*	4 (330-540 AD) 3 (260-540 AD)	house house	posthole posthole
Finstad nedre	137/1	settlement	9	7	1	1	448	445**	2	1	Cerealia Gallium aparine	2 (425-540 AD) 1	two aisled house two aisled house	posthole posthole
Furuset øvre	110/1	settlement / production site	25	6	3	17	263	25 (11)	4	234	Stellaria media Corylus avellana	28 1 (120-230 AD)	wind break -	posthole burial
Kneppe	36/120	settlement	20	6	1	13	131	91 (10)	1	39	Chenopodium album Chenopodium album	5 4	house longhouse	posthole posthole
Nitberg østre	33/31	settlement	5	5	0	0	215	215	0	0	Picea abies - needle leaf Picea abies - needle leaf	44 41	smithy smithy	pit pit
Nordre Moer	54/3	settlement	15	9	4	2	1889	1870 (2)	15**	4	Cerealia Euphorbia helioscopia	3 (95-30 BC) 2	four post constr. four post constr.	posthole posthole
Skogum / Fjellek	21/1	settlement	6	5	1	0	241	240	1	0	Hordeum vulgare	1	-	cooking pit
Soltun av Pyt	39/42	settlement /prod. site	7	6	1	1	323	319	2	2	Hordeum vulgare	2	-	cultural deposit
Søndre Randem	8/3	settlement	10	6	3	1	313	309	3**	1**	Triticum aestivum Cerealia	1 1	house house	posthole posthole
Gjølsjødegården	9/77	settlement	10	3	5	2	49	28	13	8	Galium sp. Hordeum vulgare var. nudum	5 4	longhouse longhouse	posthole posthole
Glemmen / Nøkleby	202/11, 203/120	settlement	24	7	3	14	1810	864 (3)	918	28**	Cerealia Cerealia	143 256	house house	wall ditch posthole
Grinden av Gon	9/2	settlement	7	6	1	0	572	571	1	0	Cerealia	1 (400-360 BC)	house with paddock	posthole
Kolberg søndre	28/6	settlement	10	9	1	0	744	742	2	0	Hordeum vulgare Hordeum vulgare	1 1	- -	cooking pit hearth
Lekum	90/1	grave field	8	7	1	0	17	16**	1	0	Hordeum vulgare var. nudum	1 (205 BC-5 AD)	-	grave
Løken søndre	54/8	settlement	5	4	1	0	217	216	1	0	Hordeum vulgare	1 (360-100 BC)	-	burial
Melleby	45/1	settlement	17	10	2	5	196	187	4	5	Persicaria maculosa* Carex nigra*	1 1 (377-197 BC)	longhouse longhouse	postpipe postpipe
Riggsum	51/1	settlement	14	11	2	1	312	307	4	1	Hordeum vulgare var. nudum Spargula arvensis*	1 (362-178 BC) 1 (1657-1950 AD)	two aisled house longhouse	posthole posthole

Locality	Farm nr.	Site type	Total no. of taxa	No. of tree taxa	No. of cereal taxa	No. of herb taxa	Total no. of macrofossils	Qu. of tree macrofossils (of which fruit/leaf/bark)	Qu. of cereal macrofossils	Qu. of herb macrofossils	Taxa of highest abundance in a feature/ Taxa of second highest abundance in a feature	Quantity (dated)	Structure	Feature	
Molle østre & vestre	2/1, 2/2	settlement	14	6	1	7	797	781 (7)	2	14	Spergula arvensis* Stellaria media*	2 2	smithy smithy	ditch post	
Opstad søndre	2073/7, 146	settlement	5	1	4	0	1211	3 (3)	1208	0	Hordeum vulgare Triticum aestivum	371 (470-395 BC) 4 (395-370 BC)	longhouse longhouse	posthole posthole	
Rom søndre	91/2	settlement	8	2	3	6	240	2**	234**	4	Hordeum vulgare Cerealia	100 100	- -	posthole posthole	
Skjersaaker / Fagerli	40/7	settlement	38	11	2	25	485	317	4	164	Chenopodium album Stellaria media	9 9	longhouse longhouse	- -	
Skøyen	5/1	settlement	18	7	3	8	867	538 (31)	308	21	Cerealia Hordeum vulgare	73 25	house longhouse	post post	
Vister	185/2, 185/7	settlement	49	6	3	40	1228	494 (1)	465	269	Cerealia Hordeum vulgare	116 112	house house	posthole posthole	
Ysteheide	196/17	settlement	3	2	0	1	11	4** (3)	0	7	Rubus saxatilis Rubus saxatilis	3 2	two aisled house two aisled house	posthole posthole	
Sum							14457	9913	2745	560					

Table 2. Table presenting the excavated locations from Viken and the reported macrofossils from each locality organized in columns presenting the total number of taxa, divided in the number of tree taxa, cereal taxa and herb taxa from each site, as well as the total number of macrofossils and how they are distributed between trees, cereal and herbs.

** value is set to 1 where quantity was not given. Fragments are counted as 1 seed. *when more than one species have the same highest quantity in a locality the example is chosen at random. Quantity is given along with available dating results. If context material has been dated (material found together with macrofossil in the same feature), this is indicated in italics.

as useful plants and edible crops. *No. of cereal taxa* sums up all identified grains that are cultivated by humans. Note that a plant part was counted as a single taxon if it is identified at family, genus, or species levels (e.g., Salicaceae and *Populus tremula* count as 1 taxon each dependent on identification level). The total quantity of macrofossil samples in a locality is given in the column *Total no. of macrofossils* and divided into three columns: *Qu. of tree macrofossils* (of which x is fruit/leaf/bark), *Qu. of cereal macrofossils* and *Qu. of herb macrofossils*. The *Taxa with the highest / second highest abundance in a feature* are identified for all the localities in Innlandet and Viken (tab.1 and 2, respectively), and enabled comparisons between the highest quantity of macrobotanicals in the various locations.

To evaluate regional variability of macrofossil diversity between sites (the third of the areas of investigation listed in the Introduction to this paper) we used a cluster analysis based on Euclidian distances using the functions *dist* and *hclust* (method “average”) from the R stats library (Venables and Ripley 2002; Oksanen *et al.* 2008) that results in a dendrogram grouping excavation site based on macrofossil similarities (fig. 2). Sites were compared based on presence or absence of the taxa included in the dataset. The resulting groupings were color-coded and organized in a map (fig. 3). Excavation sites in the map were color-coded based on the groupings resulting from the cluster analysis. GPS coordinates for each excavation site were taken from norgeskart.no, as coordinates given in archaeological reports sometimes vary. The position of a locality was found by searching the farm number and municipality. Maps were made using QGIS3.14.16.⁶

Results

Investigations confirm that there are observable patterns in the distribution of archaeobotanical remains within site types, structures, and features from this time period and area (fig. 1). *Site types*: the majority of archaeobotanical remains in the excavations from 400 BC to AD 400 are from settlements (71%), settlements/cultivation traces (8%) and settlement/production sites (5%). *Structures*: most archaeobotanical remains are preserved from buildings (61%); over a third of the structures are not determined. *Features*: the archaeobotany distribution over features are somewhat more varied. About half of the archaeobotanical remains (53%) are found in construction elements (such as postholes, wall ditches), one-fifth (20%) in production pits (such as cooking pits, forges), and nearly 5% in deposits.

The final dataset consists of 40 localities with a total number of 25,607 macrofossil samples. Of these, 13,974 (55%) samples are tree macrofossils (charcoal, nuts, leaves etc.), 7,189 (28%) are cereal macrofossils (cultivated grains), and 4,444 (17%) are herb macrofossils (wild or cultivated herbaceous plants such as flax, goosefoots, grasses, raspberry etc.). For Viken, the data (13,218 samples)

consists of macrofossils from trees 9913 (75%), cereals 2,745 (21%), and herbs 560 (4%). The data from Innlandet (12,389 samples) consists of macrofossils from trees 4,061 (33%), cereals 4,444 (36%), and herbs 3,884 (31%). In general, these results confirm that Innlandet county has a balanced ratio between total no. of herb/tree/cereal. Viken has very little herb material represented (4%). Only sites with very few taxa (n = 3–5) lack cereals, and this is the case for two localities (Nitberg Østre and Ystehede).

The most species-rich locality in Innlandet is Åker with 50 represented taxa and in total 4,533 macrofossil samples (141 tree samples, 2,787 cereal entries, 1,605 herb samples). The least species-rich locality in Innlandet is Berg Nedre and Øvre with nine taxa and 56 macrofossil samples. In Viken, the most species-rich locality is Skjersaaker Øvre / Fagerli with 38 represented taxa and 485 macrofossil samples. The most species-sparse locality in Viken is Ystehede with only three taxa and 11 macrofossil samples.

In Innlandet, some kinds of cereals are represented with the highest abundance (at 6 out of 14 localities) and the second-highest abundance (9 out of 14 localities). Herbaceous plants are represented with the highest abundance (eight out of 14 localities) and the second-highest abundance (5 out of 14 localities). In Viken, cereals are represented with the highest abundance (16 out of 26 localities), and the second-highest abundance (8 out of 21 localities). Herbaceous plants are represented with the highest abundance (8 out of 26 localities) and the second-highest abundance (9 out of 21 localities).

The macrobotanicals found in the largest quantities per samples are *Stellaria media* (Åker, house, posthole), *Avena* (Nyhuset Haukstad, furnace), *Chenopodium album* (Valum gård, house area, posthole; Dæhlen, house area, posthole), *Hordeum vulgare var. vulgare* (Valum gård, house area, feature na.), *Hordeum vulgare* (Opstad Søndre, house, posthole; Rom Søndre, house, posthole; Vister, house, posthole), in addition to the undefined *Cerealia* (Dæhlen, house area, posthole; Glemmen Vestre/Nøkleby Vestre, house, wall ditch/ house, posthole; Rom Søndre, posthole; Skøyen, house, post; Vister, house, posthole).

The cluster analysis shows 11 groups of localities based on similar composition of taxa (fig. 2). The possible geographical patterns in the groupings of the dendrogram are visualized in a map (fig. 3) showing the localities in colours that correspond to the dendrogram cluster groups. The taxa that are common to the cluster group are displayed under the map (fig. 3); however, localities often contain more than the taxa that were common to the cluster (as seen in tables 1–2). The light blue cluster consists of 11 sites that have two taxa in common. Three trends are identified. First, most sites across the studied area have little botanical macrofossil diversity (see light brown, grey sites; fig. 3). Second, some sites are highly diverse and are concentrated in specific areas (see brown, yellow), especially in the north of the study areas.

Sites with color according to cluster




Figure 3. Map showing counties Viken and Innlandet. Sites have colors that correspond with how they cluster in the dendrogram (fig. 2). Taxa that are common to a cluster group are displayed under the map. Individual localities may hold more taxa than the taxa common to the identified cluster group.

Characteristic taxa per cluster

- Betula, Quercus, Pinus, Corylus
- Pinus, Alnus
- Cerealia, Hordeum, Betula, Pinus, Corylus, Populus, Tilia, Chenopodium, Stellaria, Prunus, Sorbus, Viola
- Hordeum, Betula, Pinus, Galium, Chenopodium, Thlaspi
- Cerealia
- Cerealia, Hordeum, Betula, Pinus, Corylus, Quercus, Fraxinus, Tilia
- Cerealia, Betula, Pinus, Sparganium, Persicaria, Chenopodium, Stellaria, Alnus, Picea
- Cerealia, Hordeum, Betula, Pinus, Galium, Rubus, Sparganium, Carex, Persicaria, Chenopodium, Stellaria

Whilst highly diverse sites are usually found in the north, the three most diverse sites (Vister, Glemmen Vestre, and Nøkleby Vestre) are found in the south of the studied region.

Discussion

The majority of archaeobotanical samples are taken from houses within settlements, which offers potential for studying human utilization of plants in these settlements. Samples from other site-types, as well as contextual samples

from the outfield of archaeological localities, would be of great interest to better understand the dynamics between infield and outfield, as well as to broaden the understanding of various use areas of plants in the past.

The results reveal a positive relation between the diversity of features in a site, the diversity of macrofossil botanical taxa, and the quantity of samples. This could indicate that human-plant interference has been diversely distributed in the different sites, and potentially that

macrobotanical traces can be taken as indicators of past activities. No significant correlation between macrofossil diversity and specific structure and features was identified. Cultivated plants defined as *Cerealium* (*Avena* sp., *Hordeum* sp., *Secale* sp., *Triticum* sp.) appear significantly less in hearths and production pits than in other types of features. This may be an indication that the processing of cereals, roasting and drying, have not taken place in production pits, and not in random hearths, but may have been organized around specific hearths in the locations, such as in Gjølsjødegården in Viken where one hearth in the north of the house has been identified as being for roasting purposes (Kile-Vesik 2016).

Macrofossil diversity correlates with the number of herb samples, but not with the general number of macrofossil samples alone. This can be illustrated by the most species-rich locality (38 taxa) in Viken, Skjersaaker Øvre/Fagerli, which has a total of 485 macrofossils, of which 4 are cereals, 164 are herbs, and 317 are trees. Opstad Søndre and Nitberg Østre in Viken have only 5 taxa. In Opstad Søndre these are spread over 1,211 macrofossils, of which 1,208 are from cereals, none are from herbaceous plants, and three are from trees. Nitberg Østre has a total of 215 macrofossils, of which none are from cereals, none from herbaceous plants, and all 215 are from trees. The more macrofossil samples we have of herbaceous plants in a locality, the higher the diversity, which is clearly illustrated with Åker, the most species-rich locality in Innlandet, with 50 taxa, spread over 4,533 macrofossils, of which 2,787 are cereals, 1,605 are herbs, and 141 are trees. The least species-rich locality in Innlandet is Berg Nedre and Øvre (9 taxa). It has a total of 56 macrofossils, of which 5 are from cereals, 12 are from herbaceous plants, and 39 are from trees. Thus, a locality can be rich in cereal or tree macrofossils, but at the same time poor in species diversity. So what does this mean?

The correlation between herb samples and species diversity may be obvious because the Nordic flora of herbaceous plants is greater and more varied than the number of cereal varieties or species of trees. However, the result is also a reminder that changes in biodiversity, human-plant interactions, and plant practices can be revealed from studying the rich material of herbaceous plants.

Sites with high numbers of macrofossils may be a sign of more cultivation or harvest activity in an area, but can also be a result of more or specific sampling during the excavation. Moreover, since charcoal and cereals are targeted for dating purposes, trees and cereals may be overrepresented in the samples. The soil quality and preservation conditions also affect the sampling results. Seeds from different species are preserved in both varying quality and in varying quantity, where the most sturdy plant parts and the species producing the toughest and/or largest seeds may be overrepresented in the samples. Some plants are more prolific seed-producers than others: for example, *Chenopodium album* produces on

average 3,000 to 70,000 seeds per plant (<https://www.cabi.org/isc/datasheet/12648>).

Some macrofossils may be over-represented in archaeobotanical reports because of the difficulty in distinguishing between modern and archaeological specimens (Mueller-Bieniek *et al.* 2018). The appearance of many species associated with “disturbed” ruderal soils has been used as indicative of human activity (Behre 1981), yet these plants could have been key foods and medicines in the past, potentially tended, managed, or even cultivated around sites (Richer and Gearey 2017). A recurrent methodological problem is mixed soil as a consequence of post-deposition taphonomic processes, such as bioturbation, erosion, natural redeposition, and subsequent cultural activities (cultural redeposition) (Høeg 1996:9–10). Bioturbation causes uncertainties about the original deposition of seeds. However, a larger quantity of macrofossils can be seen as an argument for the macrofossils being “old” and originally deposited, since there will be less chance that bioturbation has influenced a large bulk of seeds than single found seeds. Consequently, there should be a potential in discussing use areas for herbaceous plant remains found in large quantities. Some of the presented sites above have representations of bulks of macrofossils in large quantities (>50) in postholes, defined or undefined cerealium, *Chenopodium album* and *Stellaria media*. Cereals are automatically interpreted as results of human use. But what about *Chenopodium* and *Stellaria*?

The lab analyses of macrofossils from Dæhlen suggest that the area in the excavation in which the samples of *Chenopodium* sp. seeds are found had been used in the household (Hellesøe and Skogsfjord 2010a:15 and appendix). The plant remains from the location are not treated individually in the report. However, the bulk of *Chenopodium album* can be dated via contextual finds to 180–60 BC. The archaeological report from Åker describes finds of seeds of *Stellaria media* together with finds of other typical “weeds” (Pilø 1994b), and none of these are considered potential sources in the interpretation of the place. Laboratories often distinguish between recently cultivated plants (food plants such as cereals, seeds, and berries) and “weeds” (Norw. *ugress*) (such as *Chenopodiaceae* sp., *Stellaria media* and *Centaurea cyanus*). However, the term “weed” reflects a modern understanding of often useful plants (Borgen 2020). The plants we consider weeds today may have been equally important to past settlers as cultivated crops. Indeed, many of these plants have likely been introduced to the Scandinavian area because of their value as food (e.g., *Aegopodium podagraria* and *Camelina sativa*) (Elven *et al.* 2018). The archaeobotanical lab report from Åker emphasizes the use value of *Chenopodium* as food, since seeds of the plant have been found in the stomach of one of the Danish bog bodies.

Both *Stellaria* and *Chenopodium* have been used for various purposes traditionally – especially as food and fodder (Grabowski 2014:19 with references). The first written collection of plants and their uses in a Norwegian context is Gunnerus's *Flora Norvegica* from 1776/1777 (Jørgensen *et al.* 2016). Gunnerus lists many plants used as food and fodder, but while he mentions *Stellaria* for such purposes, he does not mention *Chenopodium*. The ethnobotanical survey made by Ove Arbo Høegh in 1974 mentions *Stellaria* as fodder, but also as a useful plant for medicinal purposes, as a kind of dressing for wounds and skin problems, for dyeing, and for its ability to remove smell from hands after slaughter. He mentions the use of *Chenopodium* both as food and fodder. Gunnerus and Høegh diverge in their presentation of several plants, due to their different intentions, which in turn govern their source collection. Whereas economic growth coloured Gunnerus's collection of plant knowledge, Høegh was more interested in covering social and ritual practices, as well as medicinal ones (Teixidor-Toneu *et al.* 2020).

In Furušet Øvre, Ullensaker in Viken, two seeds of *Chenopodium album* were found in a pit potentially connected to a cremation burial. The finds are too few to provide a conclusive interpretation. Still, it could be interesting to draw attention to other graves with macrobotanical finds. Barley seeds found in a burial, barn, or grave field may be traces of different functions, e.g., ritual significance, fodder, or grave gifts. Two instances show the appearance of *Hordeum vulgare* L. (barley) in graves (Nordre Moer, Ås in Viken and Lekum, Eidsberg in Viken), however with only one seed each. The low quantity cannot be taken as an identification of an intentional deposit in either instance. But seeing the finds from different locations together may add a fresh perspective to the general understanding of how the plant has been used in the particular context.

The map showing regional variance in the period 400 BC–AD 400 (fig. 3) reveals taxa combinations in different sites. The yellow-colored excavation sites on the map have more herbs and fewer trees in common. The variety of taxa may also indicate that excavations in these sites have been carried out in soil layers where there was a more open landscape or cultivated area. The yellow sites include both *Cerealia* and *Hordeum*, which are typically recognized as cultivated crop plants. They also include the trees *Betula* and *Pinus*, as well as *Galium*, *Rubus*, *Spergula*, *Carex*, *Persicaria*, *Chenopodium*, and *Stellaria*, which are typically recognized as weeds. The similarities in sharing this diverse palette of species across localities indicate that they should be treated as useful herbaceous plants. The patterns in clustered areas are, however, quite distinct in the Viken and Innlandet regions. Brown-colored sites have many different trees, some quite short-lived: *Prunus*, *Sorbus*, *Corylus* etc. This could indicate that excavations have been carried out in layers where woods were chopped down and had opened up space for shorter-living trees, which again

could be a sign of cultivation. These sites are recognized by *Cerealia* and *Hordeum*, as well as *Chenopodium*, *Stellaria*, and *Viola*. A number of trees are also identified in all these localities: *Betula*, *Pinus*, *Corylus*, *Populus*, *Tilia*, *Prunus*, and *Sorbus*. Orange-coloured sites have a collection of longer-living trees, perhaps indicating that excavation layers are from pre-cultivated areas. These localities are, however, also characterized by *Cerealia* and *Hordeum*, as well as the trees: *Betula*, *Pinus*, *Corylus*, *Quercus*, *Fraxinus*, and *Tilia*.

One site may have been used over a long period. Although houses, postholes and ditches may have been reused, the plant remains at a certain site can be used to date agricultural activity at the site (Hjelle *et al.* 2017) and possibly rituals, cooking activity, fodder storage, and foraging. The archaeobotanical remains may not only indicate the cultivation of crops, but also, through the large amount of herbaceous plant remains, provide a window into broader past activities, including animal husbandry, foraging for food, fodder, and medicine. These cultivation activities can in many cases also be traced in the excavation sites' place names, some of which have originated in the same period.

Place names may be dated based both on linguistic and on extra-linguistic criteria. For instance, the appellative *-vin* (meaning 'natural meadow, pasture') occurs in many Norwegian toponyms (e.g., Bergen, earlier *Björgvin*, 'mountain pasture') and may be dated based on a number of phonological factors. The *-vin* element (usually an ending in a place name) commonly triggers a variety of vowel assimilations (umlauts) on the place name element it is attached to, depending on the period from which the name derives. Older and younger ages of *vin*-names may therefore be distinguished based on phonological traits of the names. For example, Helleland (2003) shows that the name *Dæli* < **dal-vin* 'valley pasture' (vowel fronting **a* > *æ*) must pre-date ca. AD 600–700, when this particular assimilation, or umlaut (i-umlaut), ceased to be active. By contrast, the cognate name *Dolve*, also derived from **dal-vin*, must be somewhat younger since it lacks i-umlaut but displays u/w-umlaut (vowel rounding **a* > *o*), active ca. AD 700–800. A name without umlaut altogether, such as *Sandven*, must post-date both of these processes (ca. AD 800–1000). In this way, certain name elements have been dated to specific periods in Nordic language development. Apart from these linguistic criteria, *vin*-names are generally also lacking from newer Norwegian settlements in the western isles (ca. AD 800–1000). Together these grammatical and distributional factors suggest that *-vin* place names date roughly to the Proto-Norse and Early Old Norse periods (ca. AD. 1–1000, NSL 1997: 493–94). The oldest names in this dataset which may stem from the period under study include the appellatives *-vin*, *-angr*, and *-heimr* (typically in reduced form, e.g., *Borgen* < *borgvin* 'settlement pasture', *Skogum* < *skógheimr* 'forest home' etc.). Many of the place

names in the dataset provide indications of continual cultivational uses, such as names indicating natural pastures, for example *Skøyen* < *skadvin* (with an unclear element *skad-*) and more generally *Leikvang* < *leikvangr* ‘sports meadow’, *Bråte* < *broti* ‘trees felled in a wood and left lying on the ground’, *Åker* < *akr* ‘field for tillage’, and *Vister* < *vistir* ‘dwellings’, among others. Even though no absolute correlation can be drawn between the place names and the archaeobotanical remains in this particular study, the examples presented here may add some perspective to the potential in investigating past activity in a place using a combination of sources.

Concluding remarks

This article has investigated the distribution of archaeobotanical remains in 40 localities from the counties of Innlandet and Viken in southeastern Norway, dated to the period 400 BC–AD 400. All botanical macrofossils from these localities have been systematized using a set of parameters that makes both archaeological and botanical information visible. Macrofossils are found mostly in features and especially in construction elements. Macrofossil diversity correlates to the number of herb samples. The compilation of a larger dataset makes it easier to compare the appearance of macrofossils across sites and to consider regional variances. A regional variance between Viken and Innlandet is confirmed. The most species-rich localities are situated in Viken, in the south of the investigation area. Still, there are generally a larger number of herb macrofossils detected in Innlandet. The quantity of taxa and macrofossils that are not trees and cereals is striking. Wild plants must have been of value for some important activities being carried out in settlements and in the outfield, and they may have played cultural, social, and economic roles. Referring to wild plants as weeds in historical analyses limits the value of considering the herbaceous plants as useful plants because of attitudes held by many people today towards these plants. Their role in past societies should not be overlooked because of a contemporary view of these plants. First when such plant remains are more systematically treated in historical disciplines will it be possible to deduce more exactly how they may have been used. Place names with Proto-Norse roots offer deep historical perspectives on land uses at certain localities, where many provide direct indications of historical cultivation activity. We hope these combined results can inspire more systematic identification and investigation of both wild and cultivated plants within and across archaeological and historical disciplines.

Notes

1. The dataset is part of a database initiative in the research project *Nordic People and Plants* collecting archaeobotanical data from archaeological excavations carried

out from the Museum of Cultural History, University of Oslo, from 1932 until the present. RCN SAMKUL funded project: Nordic People and Plants. Rediscovering and Safeguarding Nordic Ethnobotanical Heritage project no. 283364.

2. The University Museums practice different routines in sampling, analyses, and storage of archaeobotanical remains. Hjelle *et al.* (2017) describe the current situation for the university museums in Stavanger and Bergen.
3. The majority of excavations managed from the Museum of Cultural History are funded through the “polluter pays principle”. When new construction projects are planned in an area of cultural heritage (i.e. often remains of human activity) and dispensation is granted from the “Heritage Protection Act”, an excavation will take place to secure the remains for research and documentation. Excavations set out to raise and answer specific questions in combination with broad data sampling.
4. The recent county borders established during the Erna Solberg-government 2017–2021, are currently under public debate, and may be dissolved in favour of the old counties.
5. Ongoing work by Steinar Solheim, Kjetil Loftsgarden, and Frode Iversen, Museum of Cultural History, University of Oslo. See also Loftsgarden and Solheim this volume.
6. QGIS Development Team, 2021. QGIS Geographic Information System. Open Source Geospatial Foundation Project. <http://qgis.osgeo.org>), and modified in Adobe Illustrator Cs6.

Bibliography

- Behre, K. E. 1981 The interpretation of anthropogenic indicators in pollen diagrams. *Pollen et Spores* 23:225–245.
- Bharucha, Z. and J. Pretty 2010 The roles and values of wild foods in agricultural systems. *Philosophical Transactions of the Royal Society B: Biological Sciences* 365(1554):2913–2926.
- Bolker, B., M. E. Brooks, C. J. Clark, S. W. Geange, J. R. Poulsen, M. H. H. Stevens, *et al.* 2009 Generalized linear mixed models: a practical guide for ecology and evolution. *Trends in Ecology & Evolution* 24:3.
- Borgen, L. 2020 *Ugress: Et vilt herbarium*. Emil Korsmos klassiske planter. Nasjonalbiblioteket, Oslo.
- Day, J. 2013 Botany meets archaeology: people and plants in the past. *Journal of Experimental Botany* 64(18):5805–5816.
- Elven R., H. Hegre, H. Solstad, O. Pedersen, P. A. Pedersen, P. A. Åsen, *et al.* 2018, 5. juni. *Aegopodium podagraria*, vurdering av økologisk risiko. *Fremmedartslista 2018. Artsdatabanken*.

- Retrieved (2020, 2. November) from <https://www.artsdatabanken.no/fab2018/N/205>.
- Fægri, K. 1944 *Studies on the Pleistocene of western Norway. III. Bømlø*. Bergen Museum Årbog 1943, Naturvitenskapelig Serie, 8, University of Bergen, Bergen.
- Grabowski, R. 2014 *Cereal Husbandry and Settlement: Expanding Archaeobotanical Perspectives on the Southern Scandinavian Iron Age*. Unpublished PhD thesis. Umeå University, Umeå.
- Helleland, B. 2003 *Stedsnavn*. Språkrådet. <https://www.sprakradet.no/Sprakarbeid/Stedsnavn/Lar-meir-om-namn/Artiklar/Stedsnavn/>.
- Hjelle, K. L., L. Prøsch Danielsen and E. Solhjell 2017 Potential and Recommendations: Agrarian Botanical Data from Western Norway. I *The Agrarian Life of the North 2000 BC-AD 1000. Studies in Rural Settlement and Farming in Norway*, F. Iversen and H. Petersson (eds.), pp. 293–342. Cappelen Damm Akademisk, Oslo
- Hjelle, K. L. 2005 Pollenanalyse – en nødvendig metode for å forstå jernalderens jordbrukslandskap. I *Fra funn til samfunn. Jernalderstudier tilegnet Bergljot Solberg på 70-årsdagen*, K.A Bergsvik and A. jr. Engevik (eds.), pp. 91–103. UBAS Universitetet i Bergen Arkeologiske Skrifter, 1 University of Bergen, Bergen.
- Høeg, H. I. 1996 *Pollenanalytiske undersøkelser i «Østerdalsområdet» med hovedvekt på Rødsmoen, Åmot, Hedmark*. Varia 39. Universitetets Oldsaksamling, Oslo.
- Høeg, H. I., K. E. Henningsmoen and R. Sørensen 2019 Utviklingen av sen-glasial og holocen vegetasjon på Sørøstlandet, presentert i et 14C-datert standard pollendiagram. *Norges Botaniske Annaler*, 103–115.
- Høeg, O. A. 1974 *Planter og tradisjon: Floraen i levende tale og tradisjon i Norge 1925-1973*. Universitetsforlaget, Oslo, Bergen, Tromsø.
- Hverven, S. 2018 *Naturfilosofi*. Dreyers forlag, Oslo.
- Jacomet, S. 2013 Archaeobotany: analyses of plant remains from waterlogged archaeological sites. I *The Oxford Handbook of Wetland Archaeology*, F. Menotti and A. O'Sullivan (eds.), pp. 497–514. Oxford University Press, Oxford
- Jørgensen, P.M., Weidemann, E., Fremstad, E. 2016 *Flora Norvegica av J.E. Gunnerus På norsk og med kommentarer*. NTNU University Museum, *Gunneria* 80:1–505.
- Karg, S. 2012 Oil-rich seeds from prehistoric contexts. *Acta Palaeobotanica* 52(1):17–24.
- Kreuz, A. and E. Schäfer 2002 A new archaeobotanical database programme. *Vegetation History and Archaeobotany* 11, 177–179.
- Lotman, J. 2008 [1996] *Kultursemiotikk*. Cappelen Forlag, Oslo.
- Mueller-Bieniek, A., J. Pyzel and M. Kapcia 2018 *Chenopodium Seeds in Open-Air Archaeological Sites – How to Not Throw the Baby Out with the Bathwater*. *Environmental Archaeology* (25)3:1–13. doi:10.1080/14614103.2018.1536500.
- Myhre, B. 2002 Landbruk, landskap og samfunn 4000 f. Kr.–800 e.Kr. I *Norges landbrukshistorie I*, B. Myhre and I. Øye, pp. 12–213. Samlaget, Oslo.
- NSL = *Norsk Stadnamleksikon* 1997, J. Sandnes and O. Stemshaug. 4th edn. Det norske samlaget, Oslo.
- Oksanen J., R. Kindt, P. Legendre, B. O'Hara, G. L. Simpson, P. Solymos, et al. 2008 *The vegan package*. Retrieved 14 April 2016. <http://vegan.r-forge.r-project.org/>.
- Pokorná, A., Dreslerová, D., Křivánková 2011 Archaeobotanical Database of the Czech Republic, an Interim Report. *Interdisciplinaria Archaeologica – Natural Sciences in Archaeology (IANSa)* 1/II, 49–53.
- Richer S. and B. Gearey 2017 The Medicine Tree: Unsettling palaeoecological perceptions of past environments and human activity. *Journal of Social Archaeology* 17(3):239–262. doi:10.1177/1469605317731013.
- Richer, S. and B. Gearey 2018 From Rackham to REVEALS: Reflections on Palaeoecological Approaches to Woodland and Trees, *Environmental Archaeology* 23(3):286–297, doi:10.1080/14614103.2017.1283765.
- RStudio Team 2020 *RStudio: Integrated Development for R*. RStudio, PBC, Boston, MA URL <http://www.rstudio.com/>.
- Sofield, C. 2017 Thresholds in the Lives of Settlements: Anglo-Saxon Placed Deposits made at Entrances and 'Liminal Times'. I *Life on the Edge: Social, Religious and Political Frontiers in Early Medieval Europe*, S. Semple, C. Orsini and S. Mui (eds.), pp. 195–210. Neue Studien zur Sachsenforschung 6, Braunschweigesches Landesmuseum with the Internationales Sachsensymposium. Braunschweigesches Landesmuseum, Wendeburg.
- Solberg, B. 2003 [2000] *Jernalderen i Norge. Ca. 500 f.Kr. til ca. 1030 e.Kr.* Cappelen Akademisk Forlag, Oslo.
- Sture, M. 2016 Plantemateriale frå jordprøver: uviss alder og ubrukt potensial? I *Tverrfaglige perspektiver* 3, W. Brun and E. S. Pedersen (eds.), *AmS-Varia* 58, 5–20. doi:10.31265/ams-varia.v0i58.189.
- Taylor, B. 2020 Plants as persons: perceptions of the natural world in the North European Mesolithic. *Time and Mind* 13(3):1–24. doi:10.1080/1751696X.2020.1815292.
- Teixidor Toneu, I., K. Kjesrud, E. Bjerke, K. A. Parekh and A. Kool 2020 From the “Norwegian Flora” (Eighteenth Century) to “Plants and Tradition” (Twentieth Century): 200 Years of Norwegian Knowledge about Wild Plants. *Economic Botany* 74: 398–410.
- Turner, N. J., Ł. J. Łuczaj, P. Migliorini, A. Pieroni, A. L. Dreon, L. E. Sacchetti, et al. 2011 Edible

- and tended wild plants, traditional ecological knowledge and agroecology. *Critical Reviews in Plant Sciences* 30(1–2):198–225.
- van der Veen, M. 2018 Archaeobotany: the archaeology of human-plant interactions. I *The Science of Roman History. Biology, Climate, and the Future of the Past*, W. Scheidel (ed.), pp. 53–95. Princeton University Press, Princeton NJ.
- Venables W.N. and B. D. Ripley 2002 *Modern applied statistics with S*. Springer, New York
- Welinder, S., E. A. Pedersen and M. Widgren 2004 *Det svenska jordbrukets historia: Jordbrukets första femtusen år*. Nordiska museet, Stiftelsen Lagersberg.
- Archaeological excavation reports, primary sources for dataset**
- Axelsen, I. and A. Sand-Eriksen 2018 Grav, kokegropfelt, dyrkningsspor og bosetningsspor, Ljøstad vestre, 293/1, Ljøstad østre, 296/1, Stange kommune, Hedmark. Unpublished report, Kulturhistorisk museum, Universitetet i Oslo, Oslo. Retrieved from <https://www.duo.uio.no/handle/10852/61327>.
- Berge, S. L. 2010 Bosetningsspor og kokegroper. Finstad Nordre, 137/1, Ski kommune, Akershus. Unpublished report, Kulturhistorisk museum, Universitetet i Oslo, Oslo. Retrieved from <https://www.duo.uio.no/handle/10852/56072>.
- Berge, S. L., K. E. Sæther and B. Gaut 2013 Bosetningsspor, Grinden av Gon, 9/2, Rygge kommune, Østfold. Unpublished report, Kulturhistorisk museum, Universitetet i Oslo, Oslo. Retrieved from <https://www.duo.uio.no/handle/10852/53881>.
- Bukkemoen, G. B. 2012 Bosetningsspor; hus, kokegroper, graver og grøfter fra eldre og yngre jernalder. Rv2 Glåmdalen, delrapport 1. Leikvang, 14/2, Sør-Odal, Hedmark. Unpublished report, Kulturhistorisk museum, Universitetet i Oslo, Oslo. Retrieved from <https://www.duo.uio.no/handle/10852/56128>.
- Derrick, M., Sharpe, J.C. 2014. Bosetningsspor og gravminne. Melleby 45/1, Foss 44/1, Riggsum 51/1 og Jaren 46/1, Hobøl, Østfold. Unpublished report, Kulturhistorisk museum, Universitetet i Oslo, Oslo. Retrieved from <https://www.duo.uio.no/handle/10852/55816>.
- Eggen, I. M. 2010 Bosetningsspor, Skogum med Fjelken Øvre, 21/1, Asker, Akershus. Unpublished report, Kulturhistorisk museum, Universitetet i Oslo, Oslo. Retrieved from <https://www.duo.uio.no/handle/10852/51744>.
- Eggen, I. M. 2011 Bosetningsspor. Vevla, 26/1, Skjelve Store, 58/49, Stange Hedmark. Unpublished report, Kulturhistorisk museum, Universitetet i Oslo, Oslo. Retrieved from <https://www.duo.uio.no/handle/10852/65606>.
- Eikrem, K. 2008 Bosetningsspor. Opstad Søndre, 2073/7, 146, Sarpsborg kommune, Østfold fylke. Unpublished report, Kulturhistorisk museum, Universitetet i Oslo, Oslo. Retrieved from <https://www.duo.uio.no/handle/10852/56008>.
- Fyllingen, H. 2008 Bosetningsspor fra jernalderen. E18-Spydeberg, Molle Østre 2/1, Molle Vestre 2/2 og Skøyen 5/1, Spydeberg kommune, Østfold. Unpublished report, Kulturhistorisk museum, Universitetet i Oslo, Oslo. Retrieved from <https://www.duo.uio.no/handle/10852/56003>.
- Hellesøe, H. P. and A. Skogsfjord 2010a Boplassfunn og dyrkningsspor. Dæhlen 66/6, Gran kommune, Oppland. Unpublished report, Kulturhistorisk museum, Universitetet i Oslo, Oslo. Retrieved from <https://www.duo.uio.no/handle/10852/56158>.
- Hellesøe, H. P. and A. Skogsfjord 2010b Boplassfunn og dyrkningsspor. Musdalen, Vasserud 130/3, Berg 138/1,4 og Hong 141/1,2, Øyer kommune, Oppland. Unpublished report, Kulturhistorisk museum, Universitetet i Oslo, Oslo. Retrieved from <https://www.duo.uio.no/handle/10852/56156>.
- Kile, J. R. 2011 Aktivitets- og produksjonsspor fra jernalder og middelalder. Rv2 Glåmdalen, delrapport 2. Melstrøm Mellom, 18/9, Sør-Odal, Hedmark. Unpublished report, Kulturhistorisk museum, Universitetet i Oslo, Oslo. Retrieved from <https://www.duo.uio.no/handle/10852/56126>.
- Kile-Vesik, J. 2015 Bosättningsspor. Ystehede, 196/17, Halden, Østfold. Unpublished report, Kulturhistorisk museum, Universitetet i Oslo, Oslo. Retrieved from <https://www.duo.uio.no/handle/10852/55869>.
- Kile-Vesik, J. 2016 Bosättningsspor, Gjølsjødegården, 97/7, Marker, Østfold. Unpublished report, Kulturhistorisk museum, Universitetet i Oslo, Oslo. Retrieved from <https://www.duo.uio.no/handle/10852/49965>.
- Kile-Vesik, J. 2018 Bosetningsspor og graver. Lekum, 90/1, Eidsberg k., Østfold. Unpublished report, Kulturhistorisk museum, Universitetet i Oslo, Oslo. Retrieved from <https://www.duo.uio.no/handle/10852/65148>.
- Loktu, L. 2016 Bosetningsspor, langhus, grophus, kokegroper og fossilt dyrkingslag. FV.33 Langsletta-Totenviken kirke, Trogstad Nedre, 16/1 og Øvre, 17/1, Hammerstad Nordre, 15/2, Hjøll, 7/1, Østre Toten, Oppland. Unpublished report, Kulturhistorisk museum, Universitetet i Oslo, Oslo. Retrieved from <https://www.duo.uio.no/handle/10852/55915>.
- Martens, V. V. and A. Skogsfjord 2010 Gårdsbosetning fra eldre jernalder. Nordre Moer, 54/3, Ås kommune, Akershus. Unpublished report, Kulturhistorisk museum, Universitetet i Oslo, Oslo. Retrieved from <https://www.duo.uio.no/handle/10852/56075>.
- McGraw, J. L. 2013 Bosetningsspor. Løken Søndre, 54/8, Askim, Østfold. Unpublished report, Kulturhistorisk

- museum, Universitetet i Oslo, Oslo. Retrieved from <https://www.duo.uio.no/handle/10852/55833>.
- Mokkelbost, M. 2014 Bosetnings-, aktivitets- og dyrkningsspor. Vold, 1/3706, Hamar k., Lund søndre, 800/1, Ringsaker k. Hedmark. Unpublished report, Kulturhistorisk museum, Universitetet i Oslo, Oslo. Retrieved from <https://www.duo.uio.no/handle/10852/56129>.
- Nielsen, O. R. 1995. Bosetningsspor og jernvinne. Ny Fv116 Nyhuset-Haukestad, Løten kommune, Hedmark fylke. Unpublished report, Kulturhistorisk museum, Universitetet i Oslo, Oslo. Retrieved from <https://www.duo.uio.no/handle/10852/59192>.
- Olsen, D. E. F. 2011 Grav, tørkegropp, kokegroper og nedgravninger fra eldre jernalder. Rv2 Glåmdalen, delrapport 4, lok. 4. Fulu store, 3/8, 10, Sør-Odal, Hedmark. Unpublished report, Kulturhistorisk museum, Universitetet i Oslo, Oslo. Retrieved from <https://www.duo.uio.no/handle/10852/56124>.
- Orvik, K. 2018 Graver og bosetningsspor. Kneppe, 36/120, Nannestad k. Garder Østre, 166/21, Ullensaker k., Akershus. Unpublished report, Kulturhistorisk museum, Universitetet i Oslo, Oslo. Retrieved from <https://www.duo.uio.no/handle/10852/65147>.
- Pilø, L. 1994a Bosetningsspor. Valum, 20/1, Hamar, Hedmark. Unpublished report, Kulturhistorisk museum, Universitetet i Oslo, Oslo. Retrieved from <https://www.duo.uio.no/handle/10852/59507>.
- Pilø, L. 1994b Bosetningsspor. Åker gård 1992–1994, 7/201, Hamar kommune, Hedmark. Unpublished report, Kulturhistorisk museum, Universitetet i Oslo, Oslo. Retrieved from <https://www.duo.uio.no/handle/10852/56301>.
- Russ, H. and A. Skogsfjord 2010 Bosetningsspor/ dyrkningsspor, Hveem Nordøstre, 72/1, Østre Toten kommune, Oppland. Unpublished report, Kulturhistorisk museum, Universitetet i Oslo, Oslo. Retrieved from <https://www.duo.uio.no/handle/10852/51477>.
- Rødberg, F. H. 2014 Bosetnings- og aktivitetsspor. Bråte, 78/11, Skedsmo k, Akershus. Unpublished report, Kulturhistorisk museum, Universitetet i Oslo, Oslo. Retrieved from <https://www.duo.uio.no/handle/10852/55812>.
- Sharpe, J. C. 2014 Bosetningsspor. Kolberg søndre, 28/6, Fredrikstad kommune, Østfold. Unpublished report, Kulturhistorisk museum, Universitetet i Oslo, Oslo. Retrieved from <https://www.duo.uio.no/handle/10852/55825>.
- Skogstrand, L. 2012 Gårdsanlegg fra eldre jernalder/ bosetningsspor. Borgen, 30/1, Sørums kommun, Akershus. Unpublished report, Kulturhistorisk museum, Universitetet i Oslo, Oslo. Retrieved from <https://www.duo.uio.no/handle/10852/56083>.
- Solberg, A. 2011 Boplasser med hus fra førromersk jernalder (R1) samt folkevandringstid og middelalder (R3), Bjørnstad 195/3 (R1) Vister, 185/2,7 (R3), EIDSBERG, ØSTFOLD. Unpublished report, Kulturhistorisk museum, Universitetet i Oslo, Oslo. Retrieved from <https://www.duo.uio.no/handle/10852/53880>.
- Sæther, K. E. 2011 Bosetnings og produksjonsspor, Glemmen Vestre, Gnr 202/203, Bnr 11/120, 391, Fredrikstad kommune, Østfold fylke. Unpublished report, Kulturhistorisk museum, Universitetet i Oslo, Oslo. Retrieved from <https://www.duo.uio.no/handle/10852/51028>.
- Sæther, K. E. 2012 Boplass og kokegroppfelt fra romertid, Brustad av Bundli, 19/1 Hurdal kommune, Akershus fylke. Unpublished report, Kulturhistorisk museum, Universitetet i Oslo, Oslo. Retrieved from <https://www.duo.uio.no/handle/10852/53887>.
- Sæther, K. E. 2012 Boplass, Skjersaaker øvre/Fagerli, 40/7, Spydeberg kommune, Østfold fylke. Unpublished report, Kulturhistorisk museum, Universitetet i Oslo, Oslo. Retrieved from <https://www.duo.uio.no/handle/10852/53883>.
- Sæther, K. E. 2013 Bosetning- og produksjonsspor, Furuset øvre, 110/1, Ullensaker kommune, Akershus fylke. Unpublished report, Kulturhistorisk museum, Universitetet i Oslo, Oslo. Retrieved from <https://www.duo.uio.no/handle/10852/53890>.
- Sæther, K. E. 2016 Gropsjaktovn og kokegroper, Børstad, 4/1, Hamar Kommune, Hedmark fylke. Unpublished report, Kulturhistorisk museum, Universitetet i Oslo, Oslo. Retrieved from <https://www.duo.uio.no/handle/10852/49812>.
- Volker, D. 2008 Bosetningspor (groper og stolpehull) fra eldre jernalder. Rom Søndre, 91/2, Askim, Østfold. Unpublished report, Kulturhistorisk museum, Universitetet i Oslo, Oslo. Retrieved from <https://www.duo.uio.no/handle/10852/55994>.
- Wenn, C. C. 2012. Bosetningsspor og kulturlag fra yngre bronsealder, eldre jernalder og yngre jernalder, Soltun av Pyt, 39/42, Vestby, Akershus. Unpublished report, Kulturhistorisk museum, Universitetet i Oslo, Oslo. Retrieved from <https://www.duo.uio.no/handle/10852/53949>.
- Wenn, C. C. 2013. Bosetningsspor fra steinalder, bronsealder, eldre jernalder, vikingtid og middelalder, veifar fra nyere tid. Østre Nitberg, 33/31 og Vestre Nitberg, 34/17, Skedsmo kommune, Akershus. Unpublished report, Kulturhistorisk museum, Universitetet i Oslo, Oslo. Retrieved from <https://www.duo.uio.no/handle/10852/55839>.

